

**A NEW SPECIES OF *HOMALOTYLUS*
(HYMENOPTERA: ENCYRTIDAE) FROM MEXICO,
PARASITOID OF *AZYA ORBIGERA ORBIGERA*
(COLEOPTERA: COCCINELLIDAE)¹**

Vladimir A. Trjapitzin² and Serguei V. Triapitsyn³

ABSTRACT: A new species of the encyrtid wasp genus *Homalotylus* Mayr is described from the state of Tamaulipas in Mexico. The type series of *H. shuvakhinae* sp. n. was reared from the coccinellid *Azya orbigera orbigera* (Mulsant), a predator of the coccid *Protospulvinaria pyriformis* (Cockerell). A key to the three related species from the *flaminus* group of *Homalotylus* is provided.

Key Words: Encyrtidae, *Homalotylus*, taxonomy, *Azya orbigera orbigera*, parasitoid, Mexico.

In 2000, Elisaveta Ya. Shuvakhina reared a series of *Homalotylus* Mayr (Hymenoptera: Encyrtidae) in the garden of Hacienda Santa Engracia, an historic hotel located near Ejido Benito Juárez, Municipio Hidalgo, Tamaulipas, Mexico. The adult parasitoids emerged from larvae of the ladybird beetle *Azya orbigera orbigera* (Mulsant) (Coleoptera: Coccinellidae) feeding upon the coccid *Protospulvinaria pyriformis* (Cockerell) (Hemiptera: Sternorrhyncha: Coccidae) on an undetermined plant. These parasitoids represent a previously unknown species of *Homalotylus*, which we describe herein as *H. shuvakhinae* n. sp. This is the first known host record of a *Homalotylus* from the coccinellid genus *Azya* Mulsant.

Terms for morphological features are those of Gibson (1997). Acronyms for depositories of specimens are as follows: BMNH, The Natural History Museum, London, England, UK; EMUT, Entomological Museum, Centro de Investigación, U.A.M. Agronomía y Ciencias, Universidad Autónoma de Tamaulipas, Ciudad Victoria, Tamaulipas, Mexico; UCRC, Entomology Research Museum, University of California, Riverside, California, USA; USNM, National Museum of Natural History, Washington, D.C., USA; ZISP, Zoological Institute, Russian Academy of Sciences, St. Petersburg, Russia. An abbreviation used in the text is: F = antennal funicle segment.

Genus *Homalotylus* Mayr, 1876

Type species: *Encyrtus flaminus* Dalman, 1820; by subsequent designation by Ashmead (1900). Synonyms: *Nobrinus* Thomson, 1876; *Mendozaniella* Brèthes, 1913; *Hemaenasioidea* Girault, 1916; *Anisotylus* Timberlake, 1919; *Lepidaplycus* E. Blanchard, 1936; *Neoaenasioidea* Agarwal, 1966.

¹ Received February 25, 2003. Accepted March 13, 2004.

² Centro de Investigación, U.A.M. Agronomía y Ciencias, Universidad Autónoma de Tamaulipas, Ciudad Victoria, Tamaulipas 87149, Mexico. E-mail: vatrpjapitzin@yahoo.com.

³ Entomology Research Museum, Department of Entomology, University of California, Riverside, California 92521, U.S.A. E-mail: sergei.triapitsyn@ucr.edu.

Taxonomy. *Homalotylus* is a well-known genus and its generic diagnosis is available elsewhere (Timberlake 1919). Depending on the classification, the genus *Homalotylus* is placed either in the tribe Homalotylini, subtribe Homalotylyna (Trjapitzin 1973, 1989) or the tribe Aphycini (Anis and Hayat 1998) of the subfamily Encyrtinae. The senior author does not agree, however, with Anis and Hayat's (1998) synonymy of Homalotylini under Aphycini because these seem to be two very different evolutionary lines of Encyrtinae, infesting basically different groups of hosts: Homalotylini parasitize active larvae of Coccinellidae and Chrysopidae whereas Aphycini attack more or less sedentary Pseudococcidae, and their respective oviposition behaviors are completely different.

The new taxon described herein belongs to the *flaminus* species group of *Homalotylus* as defined by Timberlake (1919). In this group of species, the ovipositor is not exerted and hardly visible except in distorted specimens, or only slightly exerted. *Homalotylus shuvakhinae* sp. n. clearly belongs to the subgroup of the *flaminus* species group in which the head is notably higher than wide in frontal view. The new species from Mexico is closely related to *H. flaminus* (Dalman) and *H. eytelweini* (Ratzeburg) in having the ocellar triangle distinctly acute. These three species can be distinguished from each other using the following key.

Key to species of *Homalotylus* related to *H. shuvakhinae* sp. n., females.

- 1 First and third segments of metatarsus white.....*H. flaminus* (Dalman)
or All segments of metatarsus black or dusky2
- 2 Tegula with a white base. Mesotarsus white (except distal segment dusky).....
.....*H. eytelweini* (Ratzeburg)
- or Tegula entirely black. Mesotarsus with basal segment black, second to
fourth segments light, and distal segment dusky.....*H. shuvakhinae*, sp. n.

Biology. Primary parasitoids of larvae and pupae of various Coccinellidae. Trjapitzin and Ruíz Cancino (1998, 2001) indicated host associations of the two species of *Homalotylus* from Mexico, both of which are unrelated to the new taxon described in this communication.

Homalotylus flaminus in Europe and Asia parasitizes coccinellids of the tribe Scymnini (Klausnitzer & Klausnitzer 1972, Klausnitzer, 1976). According to the determined specimens in ZISP and also Noyes (2002), *H. flaminus* is known from Bulgaria, Georgia, Israel, Mongolia, Russia, Spain, Sweden, and Uzbekistan.

Homalotylus eytelweini parasitizes coccinellids of the tribes Chilocorini, Coccinellini, Hippodamiini and Psylloborini (Klausnitzer & Klausnitzer 1972, Klausnitzer, 1976). According to the determined specimens in ZISP and also Noyes (2002), it is known from many countries in the Palearctic region (from Spain to Japan) as well as from India and Thailand in the Oriental region, Republic of South Africa in the Afrotropical region, and Guatemala in the Neotropical region.

Homalotylus shuvakhinae sp. n., described below, is known only from the type locality in Tamaulipas, Mexico. Its host, *Azya orbiger a orbiger a*, belongs to the tribe Azyini of the subfamily Coccinellinae; distribution records of this species in the USA are all from southern Florida (Gordon 1985).

***Homalotylus shuvakhinae* V. Trjapitzin and S. Triapitsyn, NEW SPECIES**
(Figs. 1-3)

Diagnosis. See the key and comments above.

Female. Length 1.84-2.03 mm (holotype 1.84 mm). Color. Body black, with slight metallic shine. Frontovertex with faint bronze luster. Antenna black except F6 and clava yellowish-white (apical half of F5 sometimes whitish). Mesonotum with slight violet-bronze-greenish luster; mesopleura with similar, but fainter, luster. Tegula entirely black. Forewing with transverse dark band reaching posterior margin. Legs mostly black, including mesotibial spur (except in one paratype where it is brownish white); mesotarsus with apex of first segment more or less light, second to fourth segments light (yellowish or brownish), and fifth segment dusky.

Head about 1.2 x higher than wide. Frontoververtex narrow; vertex 1/4 to 1/5 head width. Ocelli in slightly acute triangle (somewhat less than 60°). Distance between posterior ocelli less than distance between posterior ocelli to anterior ocellus (4/5 to 5/6); distance from posterior ocellus to eye margin 2 x (or a little less) more than distance between posterior ocelli. Occipital margin slightly concave. Malar space height less than eye height (as 3-4:7). Distance between lower eye margin 3 x more than width of vertex. Inner head margin almost straight (or only slightly convex).

Antenna (Fig. 1) inserted near oral margin. Scape slender, almost 8 x as long as wide. Pedicel about 1/3 length of scape, 2 x as long as wide. F1 slightly longer than wide, about half length of pedicel; F2 and F3 similar to F1; F4 slightly wider than preceding funicle segments; F5 subquadrate; F6 a little wider than long. Clava 3-segmented, about as long as combined length of 3 preceding flagellar segments, obliquely truncate dorsally almost from the base of first claval segment.

Mesosoma. Pronotum short, 7 x wider than long medially, its posterior margin concave. Mesoscutum 1.3-1.7 x as wide as long; notauli not reaching posterior margin of mesoscutum, with apices very close to each other but not meeting. Posterolateral angle of axilla transversely truncate and divided by short, thin keel. Scutellum about as long as wide and about as long as mesoscutum. Propodeum very short medially, 4-5 x shorter than scutellum; posterior part of propodeum strongly concave medially and acute laterally (in dorsal view).

Wings not abbreviated. Forewing (Fig. 2) 2.6-2.7 x as long as its maximum width. Costal cell narrow. Venation as in Fig. 3; marginal vein about 1.5 x as long as wide; stigmal vein almost straight, not strongly widening towards its rounded apex; postmarginal vein about as long as stigmal vein, angle between them about 30°. Linea calva narrow, 7 x as long as wide, not exceeding limits of dark band, closed beneath by 6 discal setae.


Metasoma about as long as mesosoma. Ovipositor either not exerted or only slightly exerted beyond apex of gaster. Pygostyles at level of 2/7 length of gaster (from its apex).

Sculpture. Frontoververtex, mesopleura, and propodeum with microcellulate sculpture. Mesonotum minutely reticulate; scutellum microcellulate, almost matte.

Male. Unknown.

Type material: Holotype female on card, labeled: 1. "MÉXICO, Tam., Cd. Victoria; Sta. Engracia. Jardín del hotel, 14.II.2000 (E. Ya. Chouvakhina)"; 2. "Ex. *Azya orbiger a orbiger a* Mulsant en *Protopulvinaria pyriformis*"; 3. "*Homalotylus shuvakhinae* Trjapitzin & S. Triapitsyn HOLOTYPE ♀." Holotype deposited in ZISP. Paratypes: same data as holotype, 5 females on cards [BMNH, EMUT, UCRC, USNM, and ZISP] and 1 female on slide [UCRC].

Etymology. This species is named after the collector, Mrs. Elisaveta Yakovlevna Shuvakhina (Chouvakhina), the wife and mother of the senior and junior authors, respectively.



Figures 1-3. *Homalotylus shuvakhinae*, new species (female). (1) Antenna; (2) Forewing; (3) Forewing venation.

ACKNOWLEDGMENTS

We thank Elisaveta Ya. Shuvakhina (Russian Entomological Society, Moscow, Russia) for collecting the specimens, Larry G. Bezark (California Department of Food and Agriculture, Sacramento, California, USA) for identifying the host coccinellid, and also Valentin P. Semjanov (ZISP) and John S. Noyes (BMNH) for valuable assistance. Douglas Yanega (UCRC) and Svetlana N. Myartseva (EMUT) critically reviewed the manuscript prior to its submission. Vladimir V. Berezovskiy (UCRC) dissected and slide-mounted in Canada balsam one of the paratypes of the new species and Jung-Wook Kim (UCRC) made digital photographs from that slide using the AutoMontage system.

LITERATURE CITED

- Anis, S. B. and M. Hayat. 1998. The Indian species of *Homalotylus* (Hymenoptera: Encyrtidae). *Oriental Insects* 32:191-218.
- Ashmead, W. H. 1900. On the genera of chalcid-flies belonging to the subfamily Encyrtinae. *Proceedings of the U. S. National Museum* 22 (1202):323-412.
- Gibson, G. A. P. 1997. Chapter 2. Morphology and terminology, pp. 16-44. In, G.A.P. Gibson, J.T. Huber and J. B. Woolley (Editors). *Annotated keys to the genera of Nearctic Chalcidoidea* (Hymenoptera). NRC Research Press. Ottawa, Ontario, Canada. 794 pp.
- Gordon, R. D. 1985. The Coccinellidae (Coleoptera) of America north of Mexico. *Journal of the New York Entomological Society* 93 (1):1-912.
- Klausnitzer, B. 1976. Katalog der Entomoparasiten der mitteleuropäischen Coccinellidae (Col.). *Studia Entomologica Forestalia* (Praha) 2(7):121-130.
- Klausnitzer, B. and H. Klausnitzer. 1972. Marienkäfer (Coccinellidae). Die Neue Brehm-Bücherei. A. Ziemsen Verlag, Wittenberg Lutherstadt. 88 pp.
- Noyes, J. S. 2002. Interactive catalogue of world Chalcidoidea 2001. The Natural History Museum, Taxapad 2002, CD-ROM.
- Timberlake, P. H. 1919. Revision of the parasitic chalcidoid flies of the genera *Homalotylus* Mayr and *Isodromus* Howard, with descriptions of two closely related genera. *Proceedings of the U. S. National Museum* 56 (2293):133-194.
- Trjapitzin, V. A. 1973. [Classification of the parasitic Hymenoptera of the family Encyrtidae (Chalcidoidea). Part II. Subfamily Encyrtinae Walker, 1837]. *Entomologicheskoye Obozreniye* 52(2):416-429. [In Russian]. English translation: *Entomological Review* 52 (2): 287-295.
- Trjapitzin, V. A. 1989. [Parasitic Hymenoptera of the fam. Encyrtidae of Palaearctics]. Nauka, Leningrad Division, Leningrad. 488 pp. [In Russian].
- Trjapitzin, V. A. and E. Ruíz Cancino. 1998. *Homalotylus terminalis* (Say) (Hymenoptera: Chalcidoidea: Encyrtidae), un parasitoide de coccinélidos (Coleoptera: Coccinellidae) en el Estado de Morelos, México. *CEIBA* 38:157-160.
- Trjapitzin, V. A. and E. Ruíz Cancino. 2001. *Homalotylus cockerelli* Timberlake (Hymenoptera: Encyrtidae) in México. *Southwestern Entomologist* 26(4):377-378.